WESLEY	Sepsis Alert Checklist Time Zero	Date		
	Severe sepsis = 2 or more SIRS criteria, a suspected infection source, <u>and</u> signs/symptoms of new acute organ dysfu	function Time	RN initials	
Within 5 minutes	<ul> <li>For Inpatient: Call 2-3131 (WMC) or overhead page (WW) to activate the <u>Sepsis Rapid</u> <u>Response Team</u>         For Main ED: Call 2-3131 to activate <u>Sepsis Alert</u> </li> <li>Document time sepsis team arrives to patient's bedside – <u>repeat</u> call if team not at bedside within 5 minutes.</li> </ul>			
Within 15 minutes	<ul> <li>Sepsis order bundle initiated by Dr</li> <li>Lab draw for STAT Lactate level (green top tube on ice) and Blood Cultures (x2)</li> <li>Obtain antibiotics from Pyxis – notify pharmacy if antibiotics are not in Pyxis for STAT delivery</li> </ul>			
Within 60 minutes	<ul> <li>Start Broad Spectrum IV antibiotics after the cultures are drawn</li> <li>In the presence of SBP &lt;90, MAP&lt;65 or drop in SBP&gt;40 pts from last normal, OR lactate ≥4, Give a RAPID infusion bolus of 30ml/kg (normal saline or lactated ringers)</li> </ul>			
	<ul> <li>Weight in kg X 30 = ml of fluid (chart both start and stop times)</li> <li>If initial lactate level is &gt;2, draw a repeat lactate level within 4 hours of positive sepsis screen. (lactate order will be reflexed automatically)</li> <li>If hypotension persists after 30ml/kg bolus, or lactate ≥ 4 = Septic Shock Time</li> <li>Start IV vasopressors for persistent hypotension.</li> </ul>			
	Provider to perform focused exam within 4 hours of Septic Shock time			
Discussion must occur with the attending MD re: patient condition and implementation of the sepsis bundle (lactate, blood cultures and IV antibiotics)				
RN Patient barcode				
Not part of the medic	Not part of the medical record – Please return to Unit Clerk to route to Sepsis Coordinator			
Sepsis Alert Checklist.Docx/New_Oct2015				